

ARTICLE

Received 19 Jun 2012 | Accepted 26 Sep 2012 | Published 23 Oct 2012

DOI: 10.1038/ncomms2170

The earliest known stem-tetrapod from the Lower Devonian of China

Jing Lu¹, Min Zhu¹, John A. Long², Wenjin Zhao¹, Tim J. Senden³, Liantao Jia¹ & Tuo Qiao¹

Recent discoveries of advanced fish-like stem-tetrapods (for example, *Panderichthys* and *Tiktaalik*) have greatly improved our knowledge of the fin-to-limb transition. However, a paucity of fossil data from primitive finned tetrapods prevents profound understanding of the acquisition sequence of tetrapod characters. Here we report a new stem-tetrapod (*Tungsenia paradoxa* gen. et sp. nov.) from the Lower Devonian (Pragian, ~409 million years ago) of China, which extends the earliest record of tetrapods by some 10 million years. Sharing many primitive features with stem-lungfishes, the new taxon further fills in the morphological gap between tetrapods and lungfishes. The X-ray tomography study of the skull depicts the plesiomorphic condition of the brain in the tetrapods. The enlargement of the cerebral hemispheres and the possible presence of the pars tuberalis in this stem-tetrapod indicate that some important brain modifications related to terrestrial life had occurred at the beginning of the tetrapod evolution, much earlier than previously thought.

¹ Key Laboratory of Evolutionary Systematics of Vertebrates, Institute of Vertebrate Paleontology and Paleoanthropology, Chinese Academy of Sciences, PO Box 643, Beijing 100044, China. ² Natural History Museum of Los Angeles County, Los Angeles, California 90007, USA. ³ Department of Applied Mathematics, Research School of Physics and Engineering, The Australian National University, Canberra 0200, Australia. Correspondence and requests for materials should be addressed to M.Z. (email: zhumin@ivpp.ac.cn) or to J.L. (email: lujing@ivpp.ac.cn).

iving tetrapods, such as the frogs, turtles, birds and mammals, are a subgroup of the tetrapod lineage (tetrapod total group or tetrapodomorphs). The lineage also includes finned and limbed tetrapods that are more closely related to living tetrapods than to living lungfishes^{1–3}. Previously, the undisputed earliest known stem-tetrapod was *Kenichthys*, a finned member from the late Emsian (~399 million years ago (mya)) of Yunnan, China^{4,5}. By comparison, the oldest fossil record of the lungfish lineage (dipnomorphs), which forms the immediate sister group of tetrapodomorphs, can be traced back to the early Lochkovian (~415 mya)⁶. Thus, a ghost lineage of at least 16 million years existed for the tetrapod total group^{7,8}.

Here, we describe a new finned stem-tetrapod, *Tungsenia* paradoxa gen. et sp. nov., from the Lower Devonian (Pragian, ~409 mya) of Yunnan, China. The new taxon is ~10 million years older than *Kenichthys*, and shifts the beginning of stem-tetrapods back to the Pragian. The parietal shield (holotype, V10687) and two lower jaws (V10688.1-2) were originally described as an indeterminate osteolepid⁹. However, these specimens were not fully prepared and their phylogenetic position was open to question. After the discovery of complementary specimens (V15132.1-22) from the same locality, we further prepared the original material to unveil the medial aspect of mandible and hitherto unrecognised aspects of the neurocranial anatomy.

Results

Geological setting. The new material was recovered from a yellow sandstone layer of the Posongchong Formation, near the dam of the Qingmen Reservoir in the suburb of Zhaotong, northeastern Yunnan, South China (Fig. 1). The Posongchong fauna includes diverse galeaspid agnathans^{9–12}, placoderms^{13,14}, onychodonts, coelacanths and rhipidistians^{9,15–18} (Supplementary Table S1), as well as lingulids and plant remains^{15,19}.

This clastic fish-bearing stratum was originally named as the 'Cuifengshan Formation', by comparsion to the Cuifengshan Group (previously the Cuifengshan Formation) in neighbouring Qujing district, Yunnan^{20,21}. However, considering the lithology and faunal composition, as well as the overlying strata, the Devonian sediments underlying the marine Pojiao Formation in Zhaotong are more suggestive of the Posongchong Formation in Wenshan and Guannan, southeastern Yunnan than the Cuifengshan Group in Qujing. In this regard, the Posongchong Formation was proposed to represent the lowermost formation of the Devonian sequence in Zhaotong^{9,15-19}.

Systematic palaeontology

Osteichthyes Huxley, 1880 Sarcopterygii Romer, 1955 Tetrapodomorpha Ahlberg, 1991 *Tungsenia* gen. nov.

Etymology. Generic name after Liu Tungsen for his pioneering contribution to Paleoichthyology in China.

Type species. T. paradoxa sp. nov.

Diagnosis. A small cosmine-covered stem-tetrapod characterized by having a large orbital notch $\sim 40\%$ of total parietal shield length, pineal foramen at the middle level of orbits, a large pair of fangs on the parasymphysial plate, short robust olfactory tracts, developed cerebral cavity, pineal and parapineal recesses not extending anteriorly, hypophysial fossa far posterior to the optic nerve root and paired tongue-like recesses at the front of the hypophysial fossa.

Tungsenia paradoxa sp. nov.



Figure 1 | Location map and lithocolumn of *Tungsenia paradoxa* **gen. et sp. nov.** C, Carboniferous; D1b, Bianqinggou Formation; D1pj, Pojiao Formation; D1ps, Posongchong Formation; D1s, Suotoushan Formation; D2qj, Qujing Formation; D2qm, Qingmen Formation; D3, Zaige Formation; J, Jurassic; P, Permian; T, Triassic. Scale bar, above, 3 km; below left, 100 m; below right, 100 km.

Etymology. Specific name from the Latin *paradoxus*, unexpected.

Holotype. IVPP V10687, a parietal shield.

Referred material. Skulls, V15132.1-6; lower jaws, V10688.1-2, V15132.7-20; cheek bone, V15132.22.

Type locality. Zhaotong, Yunnan Province, China.

Formation and age. Posongchong Formation (Pragian, Early Devonian).

Diagnosis. As for the generic diagnosis.

Description. The parietal shield is about as wide as it is long (Fig. 2). In profile, the shield presents a protruding and slightly upturned snout, suggesting that the mouth was subterminal. The orbital notch, comprising $\sim 40\%$ of total parietal shield length, is the largest among that of stem-tetrapods. The pineal foramen is anteriorly positioned, corresponding to the midpoint of the orbits.

Tungsenia resembles *Kenichthys*^{4,5} and basal dipnomorphs (for example, *Youngolepis* and *Powichthys*)²²⁻²⁴ in the medioventrally directed snout, the infraorbital sensory canal running along the suture between the premaxillary and neighbouring bones, the broad orbital tectum and the well-developed basipterygoid processes (Fig. 2a–g). The elongated, broad parasphenoid and the paired internasal pits (Fig. 2b,f; Supplementary Fig. S3b) are shared with primitive sarcopterygian *Styloichthys* and basal dipnomorphs^{8,22,24,25}, but differing from the remainder of



Figure 2 | *Tungsenia paradoxa* gen. et sp. nov. from the Lower Devonian of China. Holotype (IVPP V10687), parietal shield in dorsal (**a**), ventral (**b**, photo; **f**, line drawing), ventrolateral (**c**, photo; **g**, line drawing) and anterior (**d**, photo; **e**, line drawing) views, and its rendered translucent showing endocranial cavity in dorsal view (**h**). (**i**) Restoration of external morphology. (**j**) Right mandible (IVPP V15132.7) in mesial view. (**k**) Restoration of mandible in mesial view. add.fo, adductor fossa; art.e, ethmoid articulation; ar.Vo, vomeral area; cav.nc, cavity of nasal capsule; c.hyp, canal for buccohypophysial duct; c.n.pro, canal for profundus branch of trigeminal nerve; cv, cranial cavity; c.v.pit, canal for pituitary vein; c.ll, canal for optic nerve; fe.exa, fenestra exonarina anterior; fe.v, fenestra ventralis; fo.gl, glenoid fossa; fo.icor, intercoronoid fossa; fo.pcor, precoronoid fossa; f.pin, pineal foramer; ioc, infraorbital canal; la.Co, tooth pavement on the lateral portion of coronoids; my?, myodome for eye muscle attachment?; n.ioc, notch for infraorbital canal; n.Pmx, notch for premaxillary; np, notochordal pit; orb, orbital notch; P, parasphenoid; p-mc, pores for mandibular canal; Prart, prearticular; pr.bp, basipterygoid process; soc, supraorbital canal; Sym, symphysial area; t.Co, coronoid tusks; t.De, dentary teeth; te.o, orbital tectum; t.Psym, parasymphysial plate tusk. Scale bar, 1mm.

the stem-tetrapods^{4,26–28}(Supplementary Fig. S4). A teardropshaped recess (Fig. 2c,g) is located anterodorsal to the optic foramen. A similar recess is also found in many finned stemtetrapods, such as *Gogonasus*, *Medoevia*, *Cladarosymblema* and *Litoptychus*, and has been interpreted as the attachment area for eye muscle^{26,28–31}.

Other features shared with stem-tetrapods include the opening for the pituitary vein at the base of the basipterygoid process and a flat parasymphysial dental plate^{1,30}. A large pair of fangs on the parasymphysial plate (Fig. 2j,k; Supplementary Fig. S3c,e) is

comparable to that in limbed stem-tetrapods (for example, *Ichthyostega* and *Acanthostega*)³², but absent in finned stemtetrapods, suggesting that the parasymphysial fang pair has evolved in parallel within the tetrapod lineage. A compound cheek plate, comprising a fused squamosal + quadratojugal + preopercular (Fig. 3a,b), is similar to that in the dipnomorph *Youngolepis* and some finned stem-tetrapods (Supplementary Fig. S5a-c)³³⁻³⁵. As inferred from the anterior overlap area on the cheek plate, the postorbital bone was proportionally large as in the rhizodont *Barameda*³⁶ (Supplementary Fig. S5d).



Figure 3 | *Tungsenia paradoxa* gen. et sp. nov. Compound cheek bone (Sq + Qj + Pop) (V15132.22) in external view (**a**, photo; **b**, line drawing). Left cleithrum in external view (**c**, photo; **d**, line drawing). (**e**) Tentative reconstruction based on available materials (in black). ar.mem, area for attachment of membraneous posterior wall of branchial cavity; juc, jugal canal; n.pf, notch for peoral fin; od.Cla, overlap area for clavicle; od.Ju, overlap area for jugal; od.Po, overlap area for postorbital; p.SQP, external pit on Sq + Qj + Pop; pl.Qj, quadratojugal pit-line; pl.Sq, squamosal pit-line. Scale bar, above, 1 mm; below, 1 cm.

A reconstruction of the skull and endocast based on highresolution computed tomography (HRCT) (Figs 2h and 4; Supplementary Movies 1 and 2) reveals an unexpected combination of brain features seen in Tungsenia. The short olfactory tracts and the anteriorly extending brain cavity recall the conditions in fossil sharks and placoderms^{37,38} (Supplementary Fig. S9). Besides, the brain and nerves exhibit a previously unrecognized mosaic of tetrapod and lungfish traits. The olfactory tracts of Tungsenia resemble those of basal dipnomorphs (for example, *Porolepis* and *Glyptolepis*)²⁵ in anteriorly diverging with an angle up to 60° . Posteriorly, the space for the hemispheres (cerebral cavity) is swollen and enlarged, comparable to that in many other stem-tetrapods^{27,39}. Although the enlargement of the hemispheres has independently evolved among different major groups such as cartilaginous fishes and ray-finned fishes, the evidence from *Tungsenia* suggests that the enlarged hemispheres in the tetrapodomorphs appeared during the incipient stage of their evolution (Fig. 5).

Immediately behind the cerebral cavity, the neuro-epiphysial recess presents laterally adjacent pineal and parapineal organs (Fig. 4a,c) as in basal dipnomorphs^{25,40}. However, unlike other rhipidistians (the lungfish plus tetrapod lineages)⁴¹, these two organs do not extend forward or form canal-like structures. The mesencephalic portion is more than half of the length of the cranial cavity, which recalls the condition in placoderms³⁸. Laterally, both trigeminal and profundus nerves emerge through foramina in the ethmosphenoid (Fig. 4a,b), like those of



Figure 4 | Digital restorations of *Tungsenia paradoxa* **gen. et sp. nov. (a-c) Digital endocasts of dorsal (a), ventral (b) and left lateral (c) views. (d,e) Braincase in right lateral (d) and posterior (e) views. Selected transverse CT scan slices through the endocranial cavity, showing the brain cavity at the level of the olfactory tract (f), the opening for pineal organ (g) and the anterior part of the hypophysial fossa (h) att.bc, attachment for basicranial muscle; c.a.ci, canal for internal carotid artery; c.a.pal, canal for palatine artery; c.sup.oph, canal for superficial ophthalmic nerve; c.v.ca, canal for anterior cerebral vein; c.IV, canal for trochlear nerve; c.V2 + V3, canal for maxillary and mandibular branch of trigeminal nerve; die, diencephalon; fo.hy, hypophysial fossa; mes, mesencephalon; olf, olfactory tract; pr.conn, processus connectens; re.pap, recess for parapineal organ; re.pin, recess for pineal organ; tel, telencephalon; ?PT, paired tongue-like recesses at the front of the hypophysial fossa, possibly housing the pars tuberalis. For other abbreviations see Fig. 2. Scale bar, 1 mm.**



Figure 5 | Patterns of brain evolution in selected jawed vertebrates. The brains (extant taxa) or endocasts (extinct taxa) are shown in lateral views. The bold green lines indicate that the enlarged hemispheres have independently evolved several times among different groups. The enlarged hemispheres had occurred at the initial stage of the tetrapod evolution. [†] represents extinct taxa. The vertebrate brain archetype is modified after Striedter⁴⁴ and the brains of living taxa are modified after Nieuwenhuys⁴⁵. cb, cerebellum; hem, hemisphere; med, medulla; rhom, rhombencephalon; spi, spinal cord. For other abbreviations see Fig. 4. Not to scale.

Youngolepis and Powichthys^{22,40}. The hypophysial fossa (Fig. 4b,c), as in Youngolepis and Powichthys, is stout and triangular. However, the fossa is placed in a far more posterior position relative to the optic nerve root. A pair of tonguelike recesses at the front of the hypophysial fossa (Fig. 4b,c) may topologically correspond to the paired neurocranial structures housing the pars tuberalis (PT) in living urodeles (Supplementary Fig. S8). Incidentally, a paired tube-like structure at the front of the hypophysial fossa in the dipnomorph Glyptolepis was identified as the 'pars tuberalis' (PT)^{25,7} However, it differs from the condition in Tungsenia and living urodeles in connecting directly with the brain cavity25, and additional investigation of the 'PT' of Glyptolepis is required. The distal part of the hypophysial fossa of Tungsenia is connected with a complex of parallel canals for blood vessels, as in Youngolepis and Powichthys^{22,40}. The hypophysial canal is funnel-shaped, extending posteriorly to set its opening on the parasphenoid far from the optic nerve exit (Fig. 4c; Supplementary Fig. S8a,b).

Discussion

A phylogenetic analysis of 38 taxa, including 21 stem-tetrapods, shows that *Tungsenia* is at the most basal position of the tetrapod stem lineage (Fig. 6; Supplementary Figs S1 and S2). At least two unambiguous synapomorphies demonstrate the stem-tetrapod affinity of *Tungsenia*: the pituitary vein opening anteroventral to the basipterygoid process, and the flat parasymphysial plate of mandible.

The basal phylogenetic position, unique character combination and earlier occurrence of *Tungsenia* have wide implications for the study of the tetrapod ancestry. First, *Tungsenia* pushes back the fossil record of tetrapods by some 10 million years into the Pragian. As a result, the first appearance of the tetrapod total group has now been drawn far closer to the estimated time of the lungfish-tetrapod split⁷.

Second, *Tungsenia* displays a mixture of endocranial features previously found either in basal dipnomorphs (for example, paired internasal pits, broad parasphenoid and exits for trigeminal and profundus nerves in the parietal shield) or in tetrapodomorphs (for example, the position of the pituitary vein opening). It further fills in the morphological gap between tetrapods and lungfishes, and unveils the evolutionary pattern of character changes during the initial diversification of stem-tetrapods. Different from the mosaic cranial features, the mandible of *Tungsenia* displays undisputed stem-tetrapod evolution, changes in mandible were faster than changes in endocranium.

Last, the HRCT study of Tungsenia provides unique insights into the incipient stage of tetrapod brain evolution. For a long time, our knowledge of the brain evolution in stem-tetrapods was limited owing to a small selection of well-studied taxa^{27,38,39,42}. Thus, the investigation of the brain in *Tungsenia*, the basalmost stem-tetrapod, is crucial for revealing the acquisition sequence of tetrapod brain features. The digital endocast of Tungsenia shows many plesiomorphies shared with stem-lungfishes and nonsarcopterygian taxa, such as placoderms and early sharks. Such arrangement indicates that the accelerated brain modifications (for example, developed olfactory bulbs, longitudinally positioned pineal and parapineal canals, and shorten mesencephalon) of stem-tetrapods had happened crownward of Tungsenia. On the other hand, the inferred enlarged hemispheres of Tungsenia indicate that the enlargement of the hemispheres had occurred at the beginning of the tetrapod evolution. In addition, although the homologous comparisons among the PT of living tetrapods, the 'PT' of Glyptolepis and the paired tongue-like recesses of Tungsenia still need additional investigations to test, the newly identified recesses in an early stem-tetrapod seem to provide a potential to recognize that the PT, a structure linked with some adaptation necessary for terrestrial living modes, had a deeper origin than previously considered.

Methods

Phylogenetic analysis. A phylogenetic analysis was performed on a matrix of 263 characters scored for 38 taxa (see Supplementary Methods). All characters were treated as unordered, and weighted equally. The resulting data matrix was subjected to the parsimony analysis in PAUP* (version 4.0b10)⁴³ using a heuristic search, with *Guiyu*, *Psarolepis* and *Achoania* (basal osteichthyans) specified as the outgroup.

The analysis generated five most parsimonious trees with 579 steps (Consistency index = 0.5250; Homoplasy index = 0.4750; Retention index = 0.7255; and Rescaled consistency index = 0.3809). All trees show *Tungsenia* as the most basal stem-tetrapod. The strict consensus tree (Supplementary Fig. S1a) highlights the poor phylogenetic resolution of the 'Osteolepiformes'.

X-ray micro-tomography. HRCT data were acquired at the Australian National University in 2007. Spatial resolution was set to $11.9 \,\mu$ m, $(1024)^3$ 16-bit voxels, 80 kV and 60 mA. Slice data derived from the scans were then analysed and



Figure 6 | Temporally calibrated cladogram of the simplified strict consensus tree. *Tungsenia*, the basalmost member of the stem-tetrapods (see Supplementary Methods), extends the fossil record of the tetrapod lineage by 10 million years. The first diversification of limbed tetrapods is positioned before the early Middle Devonian, based on the limbed tetrapod trackways from Poland⁴⁶. The lungfish lineage is in orange.

manipulated using Drishti software. The three-dimensional reconstructions were created in the Mimics (v13.1) software. The images of the reconstructions were exported from Mimics and VG Studio Max (v2.0), and finalized in Adobe Photoshop.

Nomenclatural acts. This published work and the nomenclatural acts it contains have been registered in ZooBank, the proposed online registration system for the International Code of Zoological Nomenclature (ICZN). The ZooBank LSIDs (Life Science Identifiers) can be resolved and the associated information viewed through any standard web browser by appending the LSID to the prefix 'http://zoobank.org'. The LSID for this publication is: urn:lsid:zoobank.org:act:39677617-77C8-4144-800E-59CE595E66DE8; urn:lsid:zoobank.org:pub: F80BD663-B04C-46E9-A245-1DCF0FB8383B.

References

- Ahlberg, P.E. & Johanson, Z. Osteolepiforms and the ancestry of tetrapods. Nature 395, 792–794 (1998).
- Coates, M.I., Ruta, M. & Friedman, M. Ever since Owen: changing perspectives on the early evolution of tetrapods. *Ann. Rev. Ecol. Evol. Syst.* 39, 571–592 (2008).
- Clack, J.A. Gaining Ground—The Origin and Evolution of Tetrapods (Indiana University Press, 2002).
- Zhu, M. & Ahlberg, P.E. The origin of the internal nostril of tetrapods. *Nature* 432, 94–97 (2004).
- Chang, M.-M. & Zhu, M. A new Middle Devonian osteolepidid from Qujing, Yunnan. Mem. Assoc. Australas. Palaeontol. 15, 183–198 (1993).
- Zhu, M. & Fan, J.-H. Youngolepis from the Xishancun Formation (Early Lochkovian) of Qujing, China. Geobios M. S. 19, 293–299 (1995).
- Müller, J. & Reisz, R.R. Four well-constrained calibration points from the vertebrate fossil record for molecular clock estimates. *BioEssays* 27, 1069–1075 (2005).
- Zhu, M. & Yu, X.-B. A primitive fish close to the common ancestor of tetrapods and lungfish. *Nature* 418, 767–770 (2002).
- Wang, J.-Q., Fan, J.-H. & Zhu, M. Early vertebrate fossils from the Early Devonian of Zhaotong District, northeastern Yunnan. *Vertebr. PalAsiat.* 34, 1–17 (1996).
- 10. Liu, Y.-H. Lower Devonian agnathans of Yunnan and Sichuan. Vertebr. PalAsiat. 13, 202–216 (1975).

- 11. Pan, J. & Wang, S.-T. New discoveries of polybranchiaspids from Yunnan Province. *Vertebr. PalAsiat.* **19**, 113–121 (1981).
- Wang, J.-Q. & Zhu, M. *Zhaotongaspis janvieri* gen. et sp. nov., a galeaspid from Early Devonian of Zhaotong, northeastern Yunnan. *Vertebr. PalAsiat.* 32, 231–243 (1994).
- 13. Ji, S.-A. & Pan, J. New Devonian petalichthyids (Placodermi) from eastern Yunnan, China. Professional Pap. Strat. Palaeontol. 27, 172–179 (1999).
- Dupret, V. A new wuttagoonaspid (Placodermi, Arthrodira) from the Lower Devonian of Yunnan (South China): origin, dispersal, and paleobiogeographic significance. J. Vertebr. Paleontol. 28, 12–20 (2008).
- Zhu, M. & Janvier, P. Un onychodontide (Vertebrata, Sarcopterygii) du Dévonien inférieur de Chine. *Compt. Rend. Sci., Ser. II* 319, 951–956 (1994).
- Lu, J. & Zhu, M. An Early Devonian (Pragian) sarcopterygian from Zhaotong, Yunnan, China. Vertebr. PalAsiat. 46, 161–170 (2008).
- Lu, J. & Zhu, M. An onychodont fish (Osteichthyes, Sarcopterygii) from the Early Devonian of China, and the evolution of the Onychodontiformes. *Proc. R. Soc. B* 277, 293–299 (2010).
- Zhu, M. *et al.* Earliest known coelacanth skull extends the range of anatomically modern coelacanths to the Early Devonian. *Nat. Commun.* 3, 772 (2012).
- Liao, W.-H., Xu, H.-K. & Wang, C.-Y. In Symposium on the Devonian System of South China. (ed. Inst. Geol. Min. Res.) 193–213 (Geological Press, 1978).
- Xian, S.-Y. & Zhou, X.-Y. In Symposium on the Devonian System of South China. (ed. Inst. Geol. Min. Res.) 176–183 (Geological Press, 1978).
- Yun, Y. In Symposium on the Devonian System of South China. (ed. Inst. Geol. Min. Res.) 151–166 (Geological Press, 1978).
- 22. Chang, M.-M. The Braincase of Youngolepis, a Lower Devonian Crossopterygian from Yunnan, South-western China. Thesis, Stockholm Univ. (1982).
- Jessen, H.L. In Problèmes actuels de Paleontologie: Evolution des Vertébrés. (ed. Lehman, J.P.) 213–222 (Colloques Internationaux CNRS, 1975).
- Jessen, H.L. Lower Devonian Porolepiformes from the Canadian Arctic with special reference to *Powichthys thorsteinssoni* Jessen. *Palaeontogr. A* 167, 180–214 (1980).
- Jarvik, E. Middle and Upper Devonian Porolepiformes from East Greenland with special reference to *Glyptolepis groenlandica* n. sp., and a discussion on the structure of the head in the Porolepiformes. *Meddr. Grønland* 187, 1–307 (1972).
- Long, J.A., Barwick, R.E. & Campbell, K.S.W. Osteology and functional morphology of the osteolepiform fish *Gogonasus andrewsae* Long, 1985, from

the Upper Devonian Gogo Formation, Western Australia. Recs. W. A. Mus 53, 1–89 (1997).

- Jarvik, E. Basic Structure and Evolution of Vertebrates. Vol. 1 (Academic Press, 1980).
 Lebedev, O.A. Morphology of a new osteolepidid fish from Russia. Bull. Mus.
- Nat. Hist. Nat. C 17, 287–341 (1995).
 29. Fox, R.C., Campbell, K.S.W., Barwick, R.E. & Long, J.A. A new osteolepiform fish from the Lower Carboniferous Raymond Formation, Drummond Basin, Oueensland. Mem. Oueensl. Mus. 38, 97–221 (1995).
- Coates, M.I. & Friedman, M. In Morphology, Phylogeny and Paleobiogeography of Fossil Fishes. (eds Elliott, D.K., Maisey, J.G., Yu, X.-B. & Miao, D.-S.) 389–416 (Verlag Dr. Friedrich Pfei, 2010).
- Long, J.A. A new osteolepidid fish from the Upper Devonian Gogo Formation, western Australia. Rec. W. A. Mus. 12, 361–377 (1985).
- Ahlberg, P.E. & Clack, J.A. Lower jaws, lower tetrapods—a review based on the Devonian genus Acanthostega. T. Roy. Soc. Edin. Earth Sci. 88, 11–46 (1998).
- Chang, M.-M. & Yu, X.-B. Reexamination of the relationship of Middle Devonian osteolepids—fossil characters and their interpretations. *Am. Mus. Novit.* 3189, 1–20 (1997).
- Chang, M.-M. In Early Vertebrates and Related Problems of Evolutionary Biology. (eds Chang, M.-M., Liu, Y.-H. & Zhang, G.-R.) 355–378 (Science Press, 1991).
- Fan, J.-H. A new species of *Thursius* from Wuding, Yunnan. Vertebr. PalAsiat. 30, 195–209 (1992).
- 36. Long, J.A. A new rhizodontiform fish from the Early Carboniferous of Victoria, Australia, with remarks on the phylogenetic position of the group. J. Vertebr. Paleontol. 9, 1–17 (1989).
- 37. Maisey, J.G. The braincase in Paleozoic symmoriiform and cladoselachian sharks. *Bull. Am. Mus. Nat. Hist.* **307**, 1–122 (2007).
- Stensiö, E.A. The brain and the cranial nerves in fossil lower craniate vertebrates. Skr. Norske, VidenskAkad. Oslo, Mat.-Naturv. Kl. 13, 1–120 (1963).
- Snitting, D. A redescription of the anatomy of the Late Devonian Spodichthys buetleri Jarvik, 1985 (Sarcopterygii, Tetrapodomorpha) from East Greenland. J. Vertebr. Paleontol. 28, 637–655 (2008).
- Clément, G. & Ahlberg, P.E. In Morphology, Phylogeny and Paleobiogeography of Fossil Fishes. (eds Elliott, D.K., Maisey, J.G., Yu, X.-B. & Miao, D.-S.) 363–377 (Verlag Dr. Friedrich Pfeil, 2010).
- Ahlberg, P.E. A re-examination of sarcopterygian interrelationships, with special reference to the Porolepiformes. Zool. J. Linn. Soc. 103, 241–287 (1991).

- Romer, A.S. The braincase of the Carboniferous crossopterygian *Megalichthys nitidus*. Bull. Mus. Comp. Zool. 82, 1–73 (1937).
- Swofford, D.L. PAUP*: Phylogenetic Analysis Using Parsimony (* and Other Methods), Version 4.0b 10 (Sinauer Associates (2003)).
- 44. Striedter, G.F. *Principles of Brain Evolution* (Sinauer Associates, 2004). 45. Nieuwenhuys, R. In The Central Nervous System of Vertebrates. (eds
- Nieuwenhuys, R., Donkelaar, H.J. & Nicholson, C.) 273–326 (Springer, 1998). 46. Niedźwiedzki, G., Szrek, P., Narkiewicz, K., Narkiewicz, M. & Ahlberg, P.E.
- Tetrapod trackways from the early Middle Devonian period of Poland. *Nature* **463**, 43–48 (2010).

Acknowledgements

We thank M.-M. Chang, X.-B. Yu, P. Ahlberg, B. Choo and G. Young for discussions; X.-F. Lu and C.-H. Xiong for specimen preparation; J.-L. Huang for illustrations; and B. Choo for life restoration. This work was supported by the Major State Basic Research Projects (2012CB821902) of MST of China, the National Natural Science Foundation of China (40930208 and 41102010), the Chinese Academy of Sciences (KZCX2-YW-156), State Key Laboratory of Palaeobiology and Straigraphy, and the ARC Discovery grant (DP0772138).

Author contributions

M.Z. designed the project. J.L. and M.Z. performed the research. M.Z., W.Z., L.J., J.L. and T.Q. did the field work. T.J.S. conducted HRCT. J.L., M.Z. and J.A.L. wrote the manuscript.

Additional information

Supplementary Information accompanies this paper at http://www.nature.com/ naturecommunications

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at http://npg.nature.com/ reprintsandpermissions/

How to cite this article: Lu, J. et al. The earliest known stem-tetrapod from the Lower Devonian of China. Nat. Commun. 3:1160 doi: 10.1038/ncomms2170 (2012).